



Information for
patients and
carers

**Steroids for Radiotherapy during
Brain Cancer treatment**

Overview

Radiotherapy can sometimes cause some swelling in the brain in and around the radiotherapy treatment area. In view of this steroids are usually prescribed to counteract this. The steroid we most commonly use when patients are having radiotherapy is dexamethasone. Steroids are prescribed with a reducing course, meaning the quantity of steroid that you take will reduce in time. Your steroid regime will be decided by your Neuro-Oncology Consultant to ensure that you reduce the steroid dose gradually.

Side effects of taking steroids

Side effects of steroid tablets can include:

Short term usage

- indigestion or heartburn
- increased appetite, which could lead to weight gain
- difficulty sleeping
- changes in mood and behaviour, such as feeling irritable or anxious
- high blood sugar or diabetes

Long term usage

- an increased risk of infections – especially chickenpox, shingles and measles
- weakening of the bones (osteoporosis)
- high blood pressure

- Cushing's syndrome – which can cause symptoms such as thin skin that bruises easily, a build-up of fat on the neck and shoulders and a red, puffy, rounded face
- eye conditions, such as glaucoma and cataracts
- mental health problems, such as depression or suicidal thoughts; get an urgent GP appointment or call 111 if this happens

Most side effects will pass once treatment stops.

Coping with side effects of taking steroids

The following tips may help reduce the side effects of steroid tablets:

- **take your tablets in the morning with breakfast** (although some specially coated tablets can be taken without food) – this may help prevent indigestion, heartburn and sleeping difficulties
- **eat a healthy, balanced diet and exercise regularly** – this may help prevent weight gain and osteoporosis
- **avoid close contact with people who are ill; especially people who have measles, chickenpox or shingles** – get medical advice as soon as possible if you think you may have been exposed to someone with an infection
- **ensure your vaccines are up-to-date** – but do not have any "live" vaccines, such as the shingles vaccine

We may need to reduce the dose or suggest taking the tablets less often (for example, every other day) if you're having side effects.

We may also sometimes recommend other medicines to take alongside steroids to protect you from some of the side effects. For example, Lansoprazole/Omeprazole should always be taken alongside steroids to protect the stomach until your course of steroids is complete.

Coming off steroids

Never stop taking steroids suddenly, please follow the reducing course that has been prescribed to you. If you start to experience an increase in neurological symptoms such as headaches or dizziness whilst reducing your steroids, seek medical advice as your steroid dose may need to be increased again. Sometimes patients may need to stay on a low dose of steroids for longer. This will be managed by your primary oncology team.

Your regime

Dexamethasone tablets

Start Date: _____

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Take _____ mg each morning and _____ mg each evening for _____ day(s), then

Contact details

Should you require further advice or information please contact-

Radiotherapy Reception/Transport queries (01772) 522900

Radiotherapy Appointment queries (01772) 522931

Sources of further information

www.lancsteachinghospitals.nhs.uk

www.nhs.uk

www.accessable.co.uk

All our patient information leaflets are available on our website for patients to access and download:

www.lancsteachinghospitals.nhs.uk/patient-information-leaflets

Lancashire Teaching Hospitals NHS Foundation Trust is not responsible for the content of external internet sites.

Lancashire Teaching Hospitals is a smoke-free site. Smoking is not permitted anywhere on any of our premises, either inside or outside the buildings. Our staff will ask you about your smoking status when you come to hospital and will offer you support and advice about stopping smoking this will include Nicotine Replacement Therapy to help manage your symptoms of withdrawal and the opportunity to speak to a nurse or advisor from the specialist Tobacco and Alcohol Care Team.

If you want to stop smoking, you can also contact the Quit Squad Freephone 0800 328 6297.

Please ask if you would like help in understanding this information. This information can be made available in large print and in other languages.

Department: Surgery
Division: Radiotherapy
Production date: May 2023
Review date: May 2026
Document Code and version: CA157 V1