

Information for patients and carers

Bapscarcare (Thick)

Scar Management Therapy Service



Statement

The instructions and advice given by your therapist and in this booklet aim to give you a good outcome, but this cannot be guaranteed. You need to follow instructions carefully, but there will always be a risk of complications which may affect the outcome. Your therapist will explain the risks, benefits and any (or no) alternatives before they ask for your consent. If you are unsure about any proposed treatment, please do not hesitate to ask for more information.

Scar healing

Any wound or surgery will cause a scar. A scar is part of your body's normal healing process. Sometimes this healing process causes excessive scar tissue to build up and form red and raised areas called hypertrophic or keloid scarring. The scar may also feel stuck down which stops your skin from moving freely. Scar healing can take up to 18 months and may be affected by:

- How much time has passed since your injury
- Infection
- Skin type e.g. pale or dark skin
- Size and/or depth of your wound
- General health

Please let us know if you suffer from any of the following conditions:

- Diabetes
- Peripheral vascular disease
- Heart disease
- Renal (kidney) problems
- Raynaud's disease
- Dermatological skin conditions
- Other

How does silicone work?

Exactly how silicone gel works is not fully understood. However, most of the research suggests that it works by maintaining moisture (hydration) and preventing the scar from drying out. Silicone gel sheets (Thick Bapscarcare) can also be effective at preventing hypertrophic and keloid scarring. It can:

- Soften and flatten raised scar tissue
- Reduce itching and redness
- Improve skin flexibility and movement

When and how should silicone sheets be used?

Your doctor or therapist will discuss the options available and will recommend the use of silicone if appropriate for you. Silicone gel sheets should only be applied to skin that is completely healed (no open wounds).

- Wash your hands before and after use
- Ensure your scar and the surrounding skin is clean and dry
- Cut the gel sheet to size with an overlap of 2cm to the surrounding skin
- Remove the plastic covering and apply the sticky surface of the gel sheet directly over the scar

IT IS IMPORTANT TO CHECK FOR ANY ADVERSE EFFECTS FROM USING SILICONE GEL SHEETS

For best results you should aim to build up use of the thick silicone gel to a minimum of 12 hours and a maximum of 23 hours per day, but it is important to do so gradually in order to let the skin get used to the silicone:

Day 1 and 2 keep the silicone in place for up to 2 hours. Remove and check for any signs of irritation (itching, increased redness, broken skin, a rash or excessive sweating).

Day 3 and 4 - 4 hours.

Day 5 and 6 - 6 hours.

If you experience irritation, remove the silicone gel sheet and wash as per care instructions below. Once the symptoms have settled restart the treatment but gradually build up use, increasing by one hour every day. If the irritation persists then stop using the silicone gel sheet completely and tell your therapist.

How long will I need to use the silicone gel?

Best results are usually seen after 2-4 months; however, you may need to use the silicone for up to 18 months. This is the amount of time scars normally take to fully mature. This varies by individual, and you will be reviewed on a regular basis by your therapist so that changes to your treatment can be made if necessary. The same sheet can be used for one month before it will need to be replaced.

Caring for the silicone gel sheets

Store the silicone gel sheets in a cool place away from direct heat as this will damage the silicone.

- Do not use any other creams, oil, or gels under the silicone sheet
- Clean the silicone gel sheets daily with a mild soap under running lukewarm water
- Rinse thoroughly
- Place the silicone gel sheet on a clean dry surface with the sticky surface facing upwards and allow to air dry (do not apply heat)
- Only use the gel sheet once it is completely dried

PLEASE ALSO READ THE MANUFACTURER INFORMATION LEAFLET

Precautions

Side effects are uncommon. Stop using the silicone gel sheet and contact your therapist as soon as possible if you experience:

- Sore, irritated, infected or broken skin
- Increased redness or a rash under, or around the gel sheet
- Wet or soggy skin that looks whitened and wrinkly
- Any loss of movement where a scar crosses a joint

Contact details

Should you require further advice or information please contact:

Burns and plastics therapy team: **01772 522379**

Physiotherapy main department: **01772 522876**

Additional contact details

Ward 4 at Royal Preston Hospital: **01772 522244**

Dressings/burns management clinic: **01772 523542**

Burns clerk for follow-up burns management clinic: **01772 528188**

Royal Preston Hospital switchboard: **01772 716565**

Sources of further information

www.lancsteachinghospitals.nhs.uk

www.nhs.uk

www.accessable.co.uk

www.patient.co.uk

www.lancsteachinghospitals.nhs.uk/veteran-aware

Follow us on social media **@lancshospitals**

Lancashire Teaching Hospitals NHS Foundation Trust is not responsible for the content of external internet sites.

All our patient information leaflets are available on our website for patients to access and download:

Lancashire Teaching Hospitals is a smoke-free site. Smoking is not permitted anywhere on any of our premises, either inside or outside the buildings. Our staff will ask you about your smoking status when you come to hospital and will offer you support and advice about stopping smoking this will include Nicotine Replacement Therapy to help manage your symptoms of withdrawal and the opportunity to speak to a nurse or advisor from the specialist Tobacco and Alcohol Care Team.

If you want to stop smoking, you can also contact Smokefree Lancashire on Freephone **08081962638**

Please ask a member of staff if you would like help in understanding this information.
This information can be made available in large print, audio, Braille and in other languages.

Our patient information group review our leaflets regularly, if you feel you would like to feedback on this information or join our reading group please contact on email address:
patientexperienceandinvolve@LTHTR.nhs.uk

Department: Core Therapies
Division: Diagnostics and Clinical (DCS)
Production date: January 2025
Review date: January 2028
JR 1252 v1